

BIBLIOGRAPHY

1. Cotton, R.G., Rodrigues, N.R., and Campbell, R.D. (1988). Reactivity of cytosine and thymine in single-base-pair mismatches with hydroxylamine and osmium tetroxide and its application to the study of mutations. *Proceedings of the National Academy of Sciences of the United States of America* 85: 4397-401.
2. Hattori, M., Shibata, A., Yoshioka, K., and Sakaki, Y. (1993). Orphan peak analysis: a novel method for detection of point mutations using an automated fluorescence DNA sequencer. *Genomics* 15: 415-417.
3. Lipshultz, R.J., Morris, D., Chee, M., Hubbell, E., Kozal, M.J., Shah, N., Shen, N., Yang, R., and Fodor, S.P. (1995). Using oligonucleotide probe arrays to access genetic diversity. *Biotech.* 19: 442-447.
4. Lui, Y-H., Bai, J., Zhu, X., Liang, X., Siemieniak, D., Venta, P.J., and Lubman, D.M. (1995). Rapid screening of genetic polymorphisms using buccal cell DNA with detection by Matrix - Assisted Laser Desorption/Ionization Mass Spectrometry. *Rapid Commun. Mass Spectrom.* 9: 735-743.
5. Nordhoff, E., Ingendoh, A., Cramer, R., Overberg, A., Stahl, B., Karas, M., Hillenkamp, F., and Crain, P.F. (1992). Matrix-assisted laser desorption/ionization mass spectrometry of nucleic acids with wavelengths in the ultraviolet and infrared. *Rapid Commun. Mass Spectrom.* 6: 771-776.
6. Orita, M., Iwahana, H., Kanazawa, H., Hayashi, K., and Sekiya, T. (1989). Detection of polymorphisms of human DNA by gel electrophoresis as single-strand conformation polymorphisms. *Proceedings of the National Academy of Sciences of the United States of America* 86: 2766-70.
7. Pomerantz, S.C., Kowalak, J.A. and McCloskey, J.A. (1993). Determination of oligonucleotide composition from mass spectrometrically measured molecular weight. *J. Am. Soc. Mass Spectrom.* 4: 204-209.